



**Paleornithological Research 2013**

Proceed. 8<sup>th</sup> Internat. Meeting Society of  
Avian Paleontology and Evolution

Ursula B. Göhlich & Andreas Kroh (Eds)



ZooBank LSID: urn:lsid:zoobank.org:pub:FD580E82-70FA-40F8-A733-EBB22C53E8A2

## **A new roller (Aves: Coraciiformes s. s.: Coraciidae) from the Early Miocene of the Saint-Gérand-le-Puy area, Allier, France**

CÉCILE MOURER-CHAUVIRÉ<sup>1</sup>, JEAN-BAPTISTE PEYROUSE<sup>2</sup> &  
MARGUERITE HUGUENY<sup>1</sup>

<sup>1</sup> *Université Claude Bernard-Lyon1, CNRS UMR 5276, Villeurbanne Cedex, France;*

*E-mail: cecile.mourer@univ-lyon1.fr; marguerite.hugueney@univ-lyon1.fr*

<sup>2</sup> *7, rue de la Grange Vertu, 71400 Autun, France; E-mail: jbpeyrouse@wanadoo.fr*

**Abstract** — *Miocoracias chenevali*, nov. gen., nov. spec., a new genus and species of roller (Aves: Coraciiformes s. s.: Coraciidae) is described from the Early Miocene of Saint-Gérand-le-Puy. This family had not yet been reported from the rich fossil avifauna of this area. Although the material is fragmentary it shows morphological characteristics different from those of the extinct family Geranopteridae, and can be attributed to the Recent family Coraciidae. The rodents found in the same locality make it possible to attribute it to the MN2a zone. An appendix gives an updated list of the fossil birds identified in the Saint-Gérand-le-Puy area.

**Key words:** Fossil birds, Early Miocene, Saint-Gérand-le-Puy, Mont Merle, Coraciidae, rollers


### **Introduction**

The Coraciiformes sensu stricto include three extinct families, the Primobucconidae, the Eocoraciidae, the Geranopteridae, and two Recent families, the Coraciidae and the Brachypteraciidae (MAYR 2009). The Primobucconidae are known from the Early Eocene of Europe and North America and the Eocoraciidae are known from the Middle Eocene of Messel, Germany. In addition a nearly complete skeleton of a new genus and species has been described from the Lower Eocene of the Green River Fm., Wyoming, United States (CLARKE *et al.* 2009). This new form, *Paracoracias occidentalis*, is placed in the suborder Coracii. Incidentally, the name Coracii has been applied to a clade that has been defined as follows: “the stem clade including all taxa more closely related to Coracioidea than to its nearest outgroup” (CLARKE *et al.* 2009: p. 587). This clade definition is equivalent to the

content of the Coraciiformes sensu stricto of MAYR (1998).

The Geranopteridae have been described in the Late Eocene or Oligocene of the Phosphorites du Quercy, in France, but in the new excavations they have been found only in Upper Eocene layers (MAYR & MOURER-CHAUVIRÉ 2000; MOURER-CHAUVIRÉ & SIGÉ 2006). They include the species *Geranopterus alatus* MILNE-EDWARDS, 1892, *Geranopterus milneedwardsi* MAYR & MOURER-CHAUVIRÉ, 2000, and some indeterminate Geranopteridae from the locality of Perrière (MOURER-CHAUVIRÉ & SIGÉ 2006). In the old collections from Quercy, the age of which is not accurately known, there is also a Coraciiforme s. s. incertae sedis species A, represented by an almost complete tarsometatarsus (PQ 1216) (MAYR & MOURER-CHAUVIRÉ 2000).

The species *Nupharanassa bohémica* MLIKOVSKÝ, 1999, described from the Lower Miocene locality of Dolnice, Czechia, (MN 4b), initially



**FIGURE 1.** Location map of the Mont Merle locality, between the classical localities of Saint-Gérand-le-Puy and Montaigu-le-Blin. The contour lines indicate the height above sea level.

placed in the family Jacanidae, has been transferred to the genus *Geranopterus* and has become *Geranopterus bohemicus* (MLIKOVSKÝ, 1999) (MOURER-CHAUVIRÉ 1999).

In Africa Coraciiformes of indeterminate family had been reported from the Early Pliocene of Langebaanweg, South Africa (RICH 1980). These Coraciiformes belong to the family Alcedinidae (OLSON 1994) and thus Coraciidae are still unknown at the present time in this avifauna. In Europe, the genus *Eurystomus* is known by an extinct species, *Eurystomus beremendensis* KESSLER, 2010, from the Late Pliocene of Beremend 26, Hungary (MN 16) and by *Eurystomus* sp., from the Early Pleistocene of Betfia 2 and Betfia 9, Romania (KESSLER 2010). In more recent localities the Coraciidae are represented by the Recent species, *Coracias garrulus* LINNAEUS, 1758 (TYRBERG 1998).

The Mont Merle locality (Fig. 1) is situated near the town of Saint-Gérand-le-Puy (Allier, France), an area known for its rich Agenian vertebrate fauna (VIRET 1929; CHENEVAL 1983a; HUGUENEY 1997). It is a limestone hill constituted by stromatolitic bioconstructions and marly sediments deposited in a lacustrine environment (WATTINNE *et al.* 2003). A quarry was opened here, but it is now filled in. Washing and sieving of marl on the top of this hill yielded various

fossil remains from fishes (mainly Cyprinidae), amphibians (Salamandridae), lizards (Lacertidae, Anguidae), amphisbaenians, snakes (Scolophoridians, Natricinae, Viperidae), marsupials (Didelphidae), insectivores (Soricidae, Talpidae), small carnivores, ungulates (Cainotheriidae, “Moschidae”), rodents (Sciuridae, Cricetidae, Eomyidae, Gliridae) and a lagomorph. Bird remains are not numerous and include only some crushed fossil bones of Palaelodidae, Laricolidae and a new Coraciidae.

Mont Merle is an unpublished new locality that has not been studied before. Some rodents make it possible to give it a dating. The Eomyid *Rhodanomys schlosseri* DEPÉRET & DOUXAMI, 1902, is represented by upper molars characterized by a continuous longitudinal ridge, and by their size which is included in the variation range observed at Fornant 11 (ENGESSER 1990) and at Chavroches (HUGUENEY *et al.* 2006). As the distribution of *R. schlosseri* covers only a short period, these teeth place the Mont Merle locality between the Paulhiac level (MN1) where *R. transiens*, predecessor of *R. schlosseri*, is present, and the upper part of the MN 2a level (top of the quarry of Montaigu-le-Blin and locality of La Chaux in Switzerland) where the species is not reported but is replaced by the Eomyid genus *Rittereria*.

## Material and methods

The anatomical terminology follows BAUMEL & WITMER 1993, and when necessary HOWARD 1929, and BALLMANN 1969a. The fossil specimens are deposited in the collection of the Université Claude Bernard-Lyon1 (acronym FSL for Faculté des Sciences de Lyon) and in the collection of the Muséum d'Histoire Naturelle de Lyon (acronym ML). Other acronyms are: EC (locality of Escamps); EC3 (locality of Escamps 3), EC4 (locality of Escamps 4), MN (Mammal Neogene), PQ (Phosphorites du Quercy), PRR (locality of Perrière), and STG (Saint-Gérard-le-Puy).

## Systematic Palaeontology

**Order Coraciiformes sensu stricto (see MAYR 1998)**

**Family Coraciidae BATSCH, 1788 (see DUBOIS & BOUR 2010)**

***Miocoracias* nov. gen.**

**Type species:** *Miocoracias chenevali* nov. spec.

**Diagnosis:** Distal part of tarsometatarsus showing a foramen vasculare distale situated relatively proximally, proximodistally elongate, and obliquely oriented. Trochlea metatarsi IV slightly shorter than trochlea metatarsi III, and trochlea metatarsi II slightly shorter than trochlea metatarsi IV. On the dorsal face trochlea metatarsi III raised and extended by a longitudinal ridge. Trochlea metatarsi III distally splayed. On the plantar face, opening of foramen vasculare distale elongate and narrow. Canalis interosseus distalis open on the plantar side and forming an elongate and narrow sulcus. Trochlea metatarsi II showing a narrow and projecting wing. Proximal rim of trochlea metatarsi III raised compared to the surface of the fossa supratrochlearis plantaris. In distal view trochleae arranged along a weakly arched line.

**Included species:** Type species only.

**Distribution:** Early Miocene, Agenian, MN 2a, Saint-Gérard-le-Puy area, Allier, France.

**Etymology:** Greek, *Mio*, from the Miocene, and *Coracias*, modern genus of roller.

***Miocoracias chenevali* nov. sp.**

(Fig. 2.1–2.7)

**Holotype:** FSL 444 229 right tarsometatarsus, distal part.

**Paratype:** FSL 444 320 right tibiotarsus, distal part, incompletely preserved.

**Diagnosis:** As for the genus.

**Horizon and locality:** Locality of Mont Merle, district of Saint-Gérard-le-Puy, Allier, France. Early Miocene, Agenian, Mammal Neogene Zone MN 2a.

**Referred material:** ML STG 4048 right ulna. This ulna comes from the old collections and it is not possible to know from which locality it has been gathered.

**Dimensions** (in mm): Right tarsometatarsus, holotype: Length as preserved, 9.0; Distal width, 5.8; Distal depth, 3.1; Width of trochlea metatarsi III, 2.1; Depth of trochlea metatarsi III, 2.4. Right tibiotarsus: Length as preserved, 11.4; Width of shaft at the level of apophysis interna ligamenti obliqui, 2.6; Depth of shaft at the same level, 2.9; Depth of lateral condyle, 4.5. Right ulna: Total length, 53.7; Proximal width, 5.5; Proximal depth, 3.7; Width of shaft in the middle, 2.3; Depth of shaft in the middle, 2.5; Distal width from the cranial border of condylus dorsalis to top of tuberculum carpale, 5.5; Depth of condylus dorsalis, 4.8.

**Etymology:** This species is named after Jacques CHENEVAL in recognition of his numerous and valuable studies on the avifauna of the Saint-Gérard-le-Puy area.

## Description and comparisons with the two Recent genera *Coracias* and *Eurystomus*

**Tarsometatarsus** (Fig. 2.1–2.3): On the dorsal face, the foramen vasculare distale is situated more proximally in *Miocoracias* and it is more proximodistally elongated than in the Recent genera. In *Coracias* and *Eurystomus*, the opening is dorsoplantarly orientated, while in *Miocoracias* the opening is slit-shaped, obliquely heading from the dorsolateral side to the plantaromedial side of the bone. This foramen is situated at the end of the outer extensor groove (HOWARD 1929).

This extensor groove is deeper in *Eurystomus* than in *Coracias*, and it is also very deep in *Miocoracias*. In *Miocoracias*, as in *Eurystomus* and *Coracias*, trochlea metatarsi IV is slightly shorter than trochlea metatarsi III and trochlea metatarsi II is slightly shorter than trochlea metatarsi IV. In the genus *Eurystomus* trochlea metatarsi III shows two projecting rims, separated by a well-expressed groove. This groove is fainter in the genera *Coracias* and *Miocoracias*. In *Eurystomus* and *Coracias* trochlea metatarsi III is relatively narrow, while it is slightly distally splayed in *Miocoracias*.

On the plantar face, in *Miocoracias*, the opening of the foramen vasculare distale is elongate and narrow, while it is more rounded in *Eurystomus* and *Coracias*. The canalis interosseus distalis, which runs from the foramen vasculare distale to the incisura intertrochlearis lateralis is not covered by a bony blade, and forms a narrow, elongate groove on the plantar side. This character is one of the main characteristics of the Coraciiformes s. s. Trochlea metatarsi II is more clearly separated from trochlea metatarsi III in *Miocoracias* than in *Eurystomus* and *Coracias*, it is narrower than in *Eurystomus* and it shows a wing less projecting plantarly than in *Coracias*. The proximal rim of trochlea metatarsi III is clearly raised compared to the surface of the fossa supratrochlearis plantaris, while in *Eurystomus* and *Coracias* this proximal part is not raised. Trochlea metatarsi IV shows a narrow and projecting wing in *Miocoracias* and *Eurystomus*, while this wing is less developed in *Coracias*. Fossa metatarsi I extends far distally in *Miocoracias* and *Eurystomus*, while it is situated more proximally in *Coracias*.

In distal view the trochleae are arranged on a faintly arched line as in both Recent genera.

**Tibiotarsus** (Fig. 2.4–2.5): The condylus medialis is not preserved. On the cranial face, the sulcus extensorius and the pons supratendineus are situated on the median axis of the bone, as in the Recent Coraciidae and Brachypteraciidae, while in most of the other Recent birds they are situated on the medial side of the bone. At the proximal part of the condylus lateralis, and on the lateral side of the pons supratendineus, there is a strong, point-shaped tubercle, the apophysis externa ligamenti obliqui (BALLMANN 1969a), and

on the medial side there is also a thin, elongate, and well projecting apophysis interna ligamenti obliqui.


Compared to the Recent genera *Coracias* and *Eurystomus*, the distal part of the tibiotarsus is less mediolaterally compressed, and the sulcus extensorius is shallower. The apophysis externa ligamenti obliqui is very sharp and projecting while its development is variable in the Recent forms: some specimens have an almost flat apophysis, situated in the continuation of the condylus lateralis, while some others have a sharp apophysis.

On the caudal face, the condylus medialis shows a wing which projects weakly medially, as in the genera *Eurystomus* and *Coracias*, and the condylus lateralis shows a wing less projecting laterally than in *Eurystomus*, but comparable to that of *Coracias*.

On the lateral face, the condylus lateralis is very rounded and the epicondylus lateralis is hardly indicated, as in *Eurystomus* and *Coracias*.

**Ulna** (Fig. 2.6–2.7): This ulna shows the characteristic shape of the rollers' ulnae. It is elongate and slender, with a slight curvature at its proximal part. The shaft is slightly flattened on its caudal side. Compared to the genus *Coracias*, the olecranon is less projecting proximally and wider at its basis. The cotyla dorsalis is extended distally by a long, narrow lip, clearly separated from the cotyla ventralis. This lip is more rounded in *Coracias*. The distal part of the fossil is not very well preserved, but the condylus ventralis is more projecting distally and the tuberculum carpale is sharper in *Miocoracias* than in *Coracias*. Still in *Miocoracias* there is a small depression on the proximal border of the tuberculum carpale, on its cranial side. This depression is absent in the Recent genera *Coracias* and *Eurystomus*.

The shape of the olecranon looks more similar to that of *Eurystomus*, the olecranon of which is short and wide at its basis. In *Miocoracias* the distal lip of the cotyla dorsalis is incurved and medially directed, while in *Eurystomus* it is dorsally directed. On the ventral side the tuberculum ligamenti collateralis ventralis is more projecting ventrally in *Eurystomus*. At the distal part the condylus ventralis is longer in proximodistal direction in *Eurystomus*.



**FIGURE 2.** *Miocoracias chenevali* nov. gen. et nov. spec. from the Early Miocene of Mont Merle, district of Saint-Gérard-le-Puy, Allier, France. 1–3, Right tarsometatarsus, distal part, holotype, FSL 444 229 (1: dorsal view, 2: plantar view, 3: distal view); 4–5, Right tibiotarsus, distal part, paratype, FSL 444 230 (4: cranial view, 5: lateral view); 6–7, Referred material, right ulna, ML STG 4048 (6: dorsal view, 7: Ventral view).

### Comparison with the extinct family Geranopteridae

**Tarsometatarsus:** The distal part of the tarsometatarsus is known in *Geranopterus alatus* by two specimens from the Late Eocene of Escamps, EC4 n°5 (MAYR & MOURER-CHAUVIRÉ 2000: fig. 8, Y–Z), and ES n°4. In *Geranopterus* the metatarsal trochleae are shorter and less indi-

vidualized than in *Miocoracias*. In *Miocoracias* trochlea metatarsi III is more raised on the dorsal face, and it is extended by a longitudinal ridge; the outer extensor groove is deeper. In *Geranopterus* the dorsal surface of the tarsometatarsus is flatter. On the plantar face the proximal rim of trochlea metatarsi III is raised compared to the surface of fossa supratrochlearis plantaris in *Miocoracias* while in *Geranopterus* the two ridges, lateral and

medial, of trochlea metatarsi III gradually merge into the surface of this fossa. Trochlea metatarsi III is narrower in *Geranopterus* than in *Miocoracias*. The same characteristics are found on a distal tarsometatarsus of *G. milneedwardsi*, from the Late Eocene of Perrière, n° FSL 330 856, different from *G. alatus* by its smaller size.

In their phylogenetic analysis CLARKE *et al.* (2009: p. 607) have used the character 68. Tarsometatarsus, distal interosseal canal: present, canal open on plantar side, forming a deep narrow sulcus between trochleae III and IV (0); absent (1). This open canal is coded as absent (1) in *Geranopterus* while it is coded as present (0) in *Eocoracias* and in the Recent Coraciidae and Brachypteraciidae. It is coded as (?) for the three species of Primobucconidae included in the analysis. This is clearly inaccurate because this character is an important synapomorphy of the clade including the Primobucconidae, Eocoraciidae, Geranopteridae, Coraciidae and Brachypteraciidae (MAYR *et al.* 2004; MAYR 2009). This character is not visible in *Eocoracias* and it is present in the Primobucconidae from the Eocene of France (MAYR *et al.* 2004) as well as in the Geranopteridae (MAYR & MOURER-CHAUVIRÉ 2000; MOURER-CHAUVIRÉ & SIGÉ 2006).

*Geranopterus bohemicus* looks very similar to *G. alatus* and differs from *Miocoracias* by its shorter trochleae metatarsi with narrow incisurae intertrochlearis, its trochlea metatarsi III which is not raised compared to the dorsal surface of the shaft and not distally splayed. It differs also by the dorsal surface of the shaft, which is flat, and by trochlea metatarsi IV, which is narrower. Finally MLIKOVSKÝ indicates in the diagnosis of this species “a rather smooth transition between the trochlea metatarsi tertii and fossa supratrochlearis plantaris” (MLIKOVSKÝ 1999: p. 122). This characteristic corresponds well to the genus *Geranopterus* and differs from the genus *Miocoracias*.

**Tibiotarsus:** In *Geranopterus alatus* the tibiotarsus is known by two distal extremities from the locality of Escamps, EC3 n°7 and EC4 n°3 (MAYR & MOURER-CHAUVIRÉ 2000: fig. 8, T–U), and in *G. milneedwardsi* by a distal part from the locality of Perrière, n° PRR 2621. In *G. alatus* the sulcus extensorius and the pons supratendineus are deeper inside the shaft. On the medial side the

epicondylus medialis is very projecting. In *Miocoracias* the pons supratendineus is more cranially brought forward, the apophysis externa ligamenti obliqui is more projecting and the epicondylus medialis is hardly visible.

In *G. milneedwardsi* the shaft is very narrow compared to the condyles; the sulcus extensorius is faintly indicated; both condyles, lateral and medial, are narrow, very close together, and strongly projecting cranially. The apophysis externa ligamenti obliqui is projecting. The pons supratendineus is situated between the proximal parts of the condyles. Both epicondyles, lateral and medial, are not well marked.

**Ulna:** In the genus *Geranopterus* only the distal part of the ulna is known. In this genus, as in *Miocoracias*, the tuberculum carpale is very sharp but the condylus ventralis is flatter and more proximodistally elongate in *Geranopterus* (see MAYR & MOURER-CHAUVIRÉ 2000: fig. 8, K–L).

#### Comparison with the small forms from Perrière

Two other, smaller forms, designated as Geranopteridae, genus and species indeterminate, have been reported from the late Eocene locality of Perrière, in the Quercy, level MN 17b (MOURER-CHAUVIRÉ & SIGÉ 2006). A distal part of tibiotarsus, n° FSL 367088, corresponds to a bird smaller than *G. milneedwardsi*. It differs from *Miocoracias* by the fact that the sulcus extensorius and the pons supratendineus are deeper between the condyles. A distal part of tarsometatarsus, n° FSL 367073, corresponds to a bird which was still smaller than the preceding one (MOURER-CHAUVIRÉ & SIGÉ 2006: pl. 1, i–j). In this form the shaft is very narrow, and triangular in cross section. Trochlea metatarsi IV is not preserved but trochlea metatarsi II is strongly plantarly displaced. This form differs from *Miocoracias* by its very narrow shaft and its trochlea metatarsi II plantarly displaced.

#### Comparison with a Coraciiformes s. s., incertae sedis, species A (n° PQ 1216)

This form is known by an almost complete tarsometatarsus illustrated in MAYR & MOURER-

CHAUVIRÉ (2000: fig. 9G–I). *Miocoracias* shares with this form the presence of an elongate and oblique distal vascular foramen, situated at the bottom of a deep outer extensor groove. But it differs from it because in *Miocoracias* trochlea metatarsi II is slightly shorter than trochlea metatarsi III while in PQ 1216 it is distinctly shorter. On the plantar face, unlike *Miocoracias*, trochlea metatarsi III does not have a raised rim, and its two ridges, lateral and medial, merge gradually into the surface of the fossa.

## Discussion

The Coraciiformes sensu stricto are represented at the present time by two families, one restricted to Madagascar, and the other mainly distributed in the tropical regions of the Old World (FRY 2001). As it was first indicated by HARRISON (1979) small non-passerine birds were much diversified in the early Tertiary of the Northern Hemisphere and occupied the ecological niches which are now occupied by the passerines. Among these small non-passerine birds there are stem group representatives of Coraciiformes s. s., Alcediniformes, Upupiformes, Caprimulgiformes, Apodiformes, Coliiformes, Psittaciformes, etc. Stem group representatives of Coracii are known in the Early Eocene of North America (CLARKE *et al.* 2009; KSEPKA & CLARKE 2010) where rollers are no longer present. These stem group representatives have progressively disappeared and some orders are now only represented by a few relict families.

The stem group family Geranopteridae is still present in the locality of Dolnice, in Czechia, the age of which is MN 4b (MLIKOVSKÝ 2002) and thus clearly younger than the age of the type locality of *Miocoracias*.

The presence of the genus *Miocoracias* fills a gap between the Eocene and the Recent forms. The small quantity of material found in the Saint-Gérand-le-Puy area can be explained by the fact that this fossil avifauna is mainly composed of waterbirds and only includes a small proportion of terrestrial birds.

The Saint-Gérand avifauna includes a large quantity of forms belonging to families, or subfamilies, the recent distribution of which is mainly tropical. CHENEVAL (1989) already indi-

cated this abundance in the aquatic avifauna, but the same observation can be made for the terrestrial avifauna. The taxa corresponding to a tropical climate are: Anatidae (genus *Mionetta*), Palaelodidae, Phoenicopteridae, Ciconiidae (tribe Leptoptilini), Threskiornithidae, Pelecanidae, Sagittariidae, Pteroclididae, Psittacidae, Coliidae, Trogonidae and Phoeniculidae. The presence of Coraciidae is in line with this presence of tropical taxa, whether exclusively (taxa with only tropical species, not migrating to temperate zones), or not. In the Recent Coraciidae, most of the species are migratory but they breed and winter inside the intertropical zone, or very close to it (FRY 2001). Two species only are long distance intercontinental migrants, the European Roller (*Coracias garrulus*) and the Dollarbird (*Eurystomus orientalis*). These tropical taxa retreated from the high and middle latitudes during the second half of the Miocene and the Pliocene, leaving in some cases emergent long-distance migrants (LOUCHART 2008).

## Acknowledgements

We thank M. MALHURET who allowed us access to the Mont Merle, Marion PETIT who helps us to collect fossil remains, Didier BERTHET for the loan of the material from the Muséum de Lyon, and Christine LEFÈVRE, from the Muséum national d'Histoire naturelle de Paris, for the loan of comparative material. We thank Antoine LOUCHART and Marco PAVIA for their careful comments which have improved the manuscript.

## References

- ALVARENGA, H.M.F. (1995): Um primitivo Membro da Ordem Galliformes (Aves) do Terciário Médio da Bacia de Taubaté, Estado de São Paulo, Brasil. – *Anais da Academia Brasileira de Ciências*, **67**: 33–44.
- BALLMANN, P. (1966): Die Vögel aus der altburdigalischen Spaltenfüllung von Wintershof (West) bei Eichstätt in Bayern. – unpubl. PhD thesis, Ludwig-Maximilians-University Munich, 115 pp., Munich (Ludwig-Maximilians-University).
- BALLMANN, P. (1969a): Die Vögel aus der altburdigalischen Spaltenfüllung von Wintershof (West) bei Eichstätt in Bayern. – *Zitteliana*, **1**: 5–60.

- BALLMANN, P. (1969b): Les oiseaux miocènes de La Grive-Saint-Alban (Isère). – *Geobios*, **2**: 157–204.
- BAUMEL, J.J. & WITMER, L.M. (1993): Osteologia. – In: BAUMEL, J. J. (ed.): Handbook of Avian Anatomy. Nomina Anatomica Avium, 2nd edition. – *Nuttall Ornithological Club*, **23**: 45–132.
- BOCHENSKI, Z. (1997): List of European fossil bird species. – *Acta zoologica cracoviensia*, **40/2**: 293–333.
- BRISSON, M.J. (1760): Ornithologia. – Vol. 6, 543 + xlviii + 146 + xxiii pp. Paris (Bauche).
- BRODKORB, P. (1970): Two fossil owls from the Aquitanian of France. – *Quarterly Journal of the Florida Academy of Science*, **32/2**: 159–160.
- CHENEVAL, J. (1983a): Les gisements de Saint-Gérard-Le-Puy: 150 ans de Paléontologie, 1833–1983. – *Revue scientifique du Bourbonnais*, **1983**: 98–107.
- CHENEVAL, J. (1983b): Révision du genre *Palaelodus* Milne-Edwards, 1863 (Aves, Phoenicopteriformes) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France). – *Geobios*, **16/2**: 179–191.
- CHENEVAL, J. (1984): Les oiseaux aquatiques (Gaviiformes à Ansériformes) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France): Révision systématique. – *Palaeovertebrata*, Montpellier, **14/2**: 33–115.
- CHENEVAL, J. (1989): Fossil bird study, and paleoecological and paleoenvironmental consequences: example from the Saint-Gérard-le-Puy deposits (Lower Miocene, Allier, France). – *Palaeogeography, Palaeoclimatology, Palaeoecology*, **73**: 295–309.
- CHENEVAL, J. & ESCUILLIÉ, F. (1992): New data concerning *Palaelodus ambiguus* (Aves: Phoenicopteriformes: Palaelodidae): Ecological and Evolutionary Interpretations. – *Natural History Museum of Los Angeles County, Science Series*, **36**: 209–224.
- CLARKE, J.A., KSEPKA, D.T., SMITH, N.A., & NORELL, M. (2009): Combined phylogenetic analysis of a new North American fossil species confirms widespread Eocene distribution for stem rollers (Aves, Coraciiformes). – *Zoological Journal of the Linnean Society*, **157**: 586–611.
- DE PIETRI, V. L. (2013): Interrelationships of the Threskiornithidae and the phylogenetic position of the Miocene ibis «*Plegadis*» *paganus* from the Saint-Gérard-le-Puy area in central France. – *Ibis*, **155**: 544–560.
- DE PIETRI, V.L., GÜNTERT, M., & MAYR, G. (2012): A *Haematopus*-like skull and other remains of Charadriiformes from the early Miocene of Saint-Gérard-le-Puy (Allier, France), In: WORTHY, T.H. & U.B. GÖHLICH (eds) 8<sup>th</sup> International Meeting of the Society for Avian Paleontology and Evolution, Abstracts, pp 15.
- DE PIETRI, V.L., MANEGOLD, A., COSTEUR, L., & MAYR, G. (2011a): A revision of the Lari (Aves: Charadriiformes) from the early Miocene of Saint-Gérard-le-Puy (Allier, France). – *Journal of Vertebrate Paleontology*, **31/4**: 812–818.
- DE PIETRI, V.L., MANEGOLD, A., COSTEUR, L. & MAYR, G. (2011b): A new species of woodpecker (Aves: Picidae) from the early Miocene of Saulcet (Allier, France). – *Swiss Journal of Palaeontology*, **130/2**: 307–314.
- DE PIETRI, V.L. & MAYR, G. (2012): An assessment of the diversity of early Miocene Scolopaci (Aves, Charadriiformes) from Saint-Gérard-le-Puy (Allier, France). – *Palaeontology*, **55/6**: 1177–1197.
- DEPÉRET, C. & DOUXAMI, H. (1902): Les vertébrés oligocènes de Pyrimont-Challonges (Savoie). – *Mémoires de la Société paléontologique Suisse*, **29**: 1–90.
- DUBOIS, A. & BOUR, R. (2010): The distinction between family-series and class-series nomina in zoological nomenclature, with reemphasis on the nomina created by Batsch (1788, 1789) and on the higher nomenclature of turtles. – *Bonn zoological Bulletin*, **57**: 149–171.
- ENGESSER, B. (1990): Die Eomyidae (Rodentia, Mammalia) der Molasse der Schweiz und Savoyens. Systematik und Biostratigraphie. – *Schweizerische Paläontologische Abhandlungen*, **112**: 1–144.
- FRY, C.H. (2001): Family Coraciidae (Rollers). – In: DEL HOYO, J., ELLIOTT, A. & SARGATAL, J. (eds): Handbook of the Birds of the World, vol. 6, Mousebirds to Hornbills, – pp. 342–376, Barcelona (Lynx edicions).
- GAILLARD, C. (1939): Contribution à l'étude des oiseaux fossiles. – *Archives du Muséum d'Histoire naturelle de Lyon*, **15**, mém. 2: 1–100.
- GERVAIS, P. (1848–1852): Zoologie et Paléontologie françaises (Animaux vertébrés), ou Nouvelles recherches sur les animaux vivants et fossiles de la France. – Vol. I, iv + 271 pp.; Vol. II, Explication des planches et divers mémoires relatifs aux animaux vertébrés; Vol. III, Atlas, 80 pls., Paris (Arthus Bertrand).
- GILL, F. & DONSKER, D. (eds) (2012): IOC World Bird Names (v2.11). Available at <http://www.world-birdnames.org>
- GÖHLICH, U.B. & MOURER-CHAUVIRÉ, C. (2005): Revision of the Phasianids (Aves: Galliformes) from the Lower Miocene of Saint-Gérard-le-Puy (Allier, France). – *Palaeontology*, **48/6**: 1331–1350.
- HARRISON, C.J.O. (1975): The taxonomic status of Milne-Edward's fossil Sulids. – *Bulletin of the British Ornithologists Club*, **95**: 51–54.
- HARRISON, C.J.O. (1979): Small non-passerine birds of the Lower Tertiary as exploiters of ecological niches now occupied by passerines. – *Nature*, **281**: 562–563.



- HARRISON, C.J.O. (1984): A revision of the fossil swifts (Vertebrata, Aves, Suborder Apodi), with description of three new genera and two new species. – *Mededelingen van der Werkgroepvoor Tertiaire en Kwartaire Geologie*, **21/4**: 157–177.
- HOWARD, H. (1929): The avifauna of Emeryville shell-mound. – *University of California Publications in Zoology*, **32/2**: 301–394.
- HUGUENEY, M. (1997): Biochronologie mammalienne dans le Paléogène et le Miocène inférieur du Centre de la France: Synthèse réactualisée. – In: AGUILAR, J.P., LEGENDRE, S. & MICHAUX, J. (eds): Actes du Congrès Biochrom'97, Montpellier, 14–17 Avril. – *Ecole Pratique des Hautes Etudes, Mémoires et Travaux de l'Institut de Montpellier*, **21**: 417–430.
- HUGUENEY, M., BERTHET, D., BODERGAT, A.-M., ESCUILLIÉ, F., MOURER-CHAUVIRÉ, C. & WATTINNE, A. (2003): La limite Oligocène-Miocène en Limagne : changements fauniques chez les mammifères, oiseaux et ostracodes des différents niveaux de Billy-Créchy (Allier, France). – *Geobios*, **36**: 719–731.
- HUGUENEY, M., BERTHET, D., ESCUILLIÉ, F. & RIVAL, J. (2006): Eomyids (Rodentia, Mammalia) in the St-Gérard-le-Puy area (Allier, France; MN2a). – *Beiträge zur Paläontologie*, **30**: 205–222.
- KESSLER, J. (=E.) (2010): Új eredmények a Kárpát-medence neogén és negyedidőszaki madárvilágához III. [New results with regard to the Neogene and Quaternary Avifauna of the Carpathian Basin, part III]. – *Földtany Közlöny*, **140/1**: 53–72.
- KSEPKA, D.T. & CLARKE, J.A. (2010): *Primobucco mcgrewi* (Aves: Coracii) from the Eocene Green River Formation: New anatomical data from the earliest constrained record of stem rollers. – *Journal of Vertebrate Paleontology*, **30/1**: 215–225.
- LAMBRECHT, K. (1931): *Cygnopterus* und *Cygnavus*, zwei fossile Schwäne aus dem Tertiär Europas. – *Bulletin du Musée royal d'Histoire naturelle de Belgique*, **7/31**: 1–6.
- LACÉPÈDE, B.G.E. de (1799): Tableaux méthodiques des mammifères et des oiseaux. Première édition, 3ème partie, Tableau des sous-classes, divisions, sous-divisions, ordres et genres des oiseaux. – 20 pp. Paris (Plassan).
- LAMBRECHT, K. (1933): Handbuch der Palaeornithologie. – 1024 pp. Berlin (Gebrüder Borntraeger).
- LINNAEUS, C. (1758): Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus differentiis, synonymis, locis, I. – 824 pp. Stockholm. (Laurentii Salvii)
- LIVEZEY, B.C. & MARTIN, L.D. (1988): The systematic position of the Miocene anatid *Anas* [?] *blanchardi* Milne-Edwards. – *Journal of Vertebrate Paleontology*, **8/2**: 196–211.
- LOUCHART, A. (2008): Emergence of long distance bird migrations: a new model integrating global climate changes. – *Naturwissenschaften*, **95**: 1109–1119.
- LYDEKKER, R. (1891a): Catalogue of the Fossil Birds. – 368 pp. London (British Museum).
- LYDEKKER, R. (1891b): On remains of a large stork from the Allier Miocene. – *Proceedings of the Zoological Society of London* **29**: 476–479.
- MAYR, G. (1998): «Coraciiforme» und «Piciiforme» Kleinvögel aus dem Mittel-Eozän der Grube Messel (Hessen, Deutschland). – *Courier Forschungsinstitut Senckenberg*, **205**: 1–101.
- MAYR, G. (2009): Paleogene Fossil Birds. – xiii + 262 pp. Berlin, Heidelberg (Springer-Verlag).
- MAYR, G. & MOURER-CHAUVIRÉ, C. (2000): Rollers (Aves: Coraciiformes s. s.) from the Middle Eocene of Messel (Germany) and the Upper Eocene of the Quercy (France). – *Journal of Vertebrate Paleontology*, **20/3**: 533–546.
- MAYR, G., MOURER-CHAUVIRÉ, C. & WEIDIG, I. (2004): Osteology and systematic position of the Eocene Primobucconidae (Aves, Coraciiformes *sensu stricto*), with first records from Europe. – *Journal of Systematic Palaeontology*, **2/1**: 1–12.
- MAYR, G. & SMITH, T. (2012): A fossil albatross from the Early Oligocene of the North Sea Basin. – *The Auk*, **129/1**: 87–95.
- MILLER, A.H. (1944): An avifauna from the Lower Miocene of South Dakota. – *University of California Publications, Bulletin of the Department of Geological Sciences*, **27**: 85–100.
- MILNE-EDWARDS, A. (1863): Mémoire sur la distribution géologique des oiseaux fossiles et description de quelques espèces nouvelles. – *Annales des Sciences Naturelles* (4), **20**: 132–176.
- MILNE-EDWARDS, A. (1867–1868): Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France, I. – 472 pp., Atlas I: pls 1–96, Paris (Victor Masson et Fils).
- MILNE-EDWARDS, A. (1869–1871): Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France, II. – 627 pp., Atlas II: pls 97–200, Paris (Librairie de G. Masson).
- MILNE-EDWARDS, A. (1892): Sur les oiseaux fossiles des dépôts éocènes de phosphate de chaux du Sud de la France. – *Compte-Rendu du Second Congrès Ornithologique International, Budapest*: 60–80.
- MLIKOVSKÝ, J. (1998): Two new owls (Aves: Strigidae) from the early Miocene of the Czech Republic, with comments on the fossil history of the subfamily Striginae. – *Buteo*, **10**: 5–21.
- MLIKOVSKÝ, J. (1999): A new jacana (Aves: Jacanidae) from the Early Miocene of the Czech Republic. – *Comptes Rendus de l'Académie des Sciences de Paris*, sér. IIa, **328**: 121–123.
- MLIKOVSKÝ, J. (2002): Cenozoic Birds of the World. Part 1: Europe. – 406 pp. Praha (Ninox Press).

- MLIKOVSKÝ, J. & GÖHLICH, U.B. (2000): A new wood-hoopoe (Aves: Phoeniculidae) from the early Miocene of Germany and France. – *Acta Societatis Zoologicae Bohemicae*, **64**: 419–424.
- MOURER-CHAUVIRÉ, C. (1993): Les Gangas (Aves, Columbiformes, Pteroclididae) du Paléogène et du Miocène inférieur de France. – *Palaeovertebrata*, **22/2–3**: 73–98.
- MOURER-CHAUVIRÉ, C. (1995): Dynamics of the Avifauna during the Paleogene and the Early Neogene of France. Settling of the Recent fauna. – *Acta zoologica cracoviensia*, **38/3**: 325–342.
- MOURER-CHAUVIRÉ, C. (1999): Position systématique de *Nupharanassa bohémica* Mlíkovský, 1999. – *Comptes Rendus de l'Académie des Sciences de Paris*, sér. IIA, **329**: 149–152.
- MOURER-CHAUVIRÉ, C. (2000): A new species of *Ameripodius* (Aves: Galliformes: Quercymegapodiidae) from the Lower Miocene of France. – *Palaeontology*, **43/3**: 481–493.
- MOURER-CHAUVIRÉ, C. & SIGÉ, B. (2006): Une nouvelle espèce de *Jungornis* (Aves, Apodiformes) et de nouvelles formes de Coraciiformes s. s. dans l'Eocène supérieur du Quercy. Journées Bernard Gèze. – *Strata*, sér. 1, **13**: 151–159.
- OLSON, S.L. (1994): Early Pliocene grebes, button-quail and kingfishers from south-western Cape Province, South Africa (Aves: Podicipedidae, Turnicidae, Halcyornithidae). – *Annals of the South African Museum*, **104/4**: 49–61.
- PATON, T.A., BAKER, A.J., GROTH, J.G. & BARROW-CLOUGH, G.F. (2003): RAG-1 sequences resolve phylogenetic relationships within Charadriiform birds. – *Molecular Phylogenetics and Evolution*, **29**: 268–278.
- PORTIS, A. (1884): Contribuzione alla ornithologia Italiana. – *Estr. dalle Memorie della Reale Accademia delle Scienze di Torino*, (2) **36**: 2–26.
- RICH, P.V. (1980): Preliminary report on the fossil avian remains from the late Tertiary sediments at Langebaanweg (Cape Province), South Africa. – *South African Journal of Science*, **76**: 166–170.
- STEJNEGER, L. (1885): Birds. – In: KINGSLEY J. S. (ed.): The standard Natural History. Vol. 4, vii + 558 pp. Boston (S. E. Cassino).
- TYRBERG, T. (1998): Pleistocene birds of the Palearctic: a Catalogue. – *Publications of the Nuttall Ornithological Club*, **27**: 1–720.
- VIRET, J. (1929): Les faunes de mammifères de l'Oligocène de la Limagne bourbonnaise. – *Annales de l'Université de Lyon, N. S.*, **1**: 1–328.
- WATTINNE, A., VENNIN, E. & DE WEVER, P. (2003): Evolution of carbonated, lacustrine environment, with stromatolites: a paleoecological approach (quarry of Montaigu-le-Blin, Limagne graben, Allier, France). – *Bulletin de la Société géologique de France*, **174/3**: 243–260.
- WETMORE, A. (1958): Miscellaneous notes on fossil birds. – *Smithsonian Miscellaneous Collections*, **135/8**: 11 pp.

## Appendix

Updated list of the avian species reported in the whole of the localities known as “Saint-Gérard-le-Puy”. The material from the old collections comes from different quarries and its age is not accurately known; this age can be MN 1, MN 1/2, or MN 2a (HUGUENEY 1997). However the largest part of the quarries has yielded fossil material corresponding to the MN 2a zone. This list is founded upon the following works: BALLMANN 1969b; CHENEVAL 1983b, 1984; CHENEVAL & ESCUILLIÉ 1992; MOURER-CHAUVIRÉ 1995; BOCHEŃSKI 1997; MLIKOVSKÝ 1998; MOURER-CHAUVIRÉ 2000; MLIKOVSKÝ & GÖHLICH 2000; MLIKOVSKÝ 2002; HUGUENEY *et al.* 2003; GÖHLICH & MOURER-CHAUVIRÉ 2005; DE PIETRI *et al.* 2011a, 2011b, 2012; MAYR & SMITH 2012; DE PIETRI & MAYR 2012; DE PIETRI 2013. The systematic order follows the list of the International Ornithologists' Union (GILL & DONSKER 2012). Some modifications brought by MLIKOVSKÝ (2002) have not been integrated because they do not seem well-founded.

### O. Galliformes

- F. Quercymegapodiidae MOURER-CHAUVIRÉ, 1992  
 Genus *Ameripodius* ALVARENGA, 1995  
*Ameripodius alexis* MOURER-CHAUVIRÉ, 2000
- F. Phasianidae HORSFIELD, 1821  
 Genus *Palaeortyx* MILNE-EDWARDS, 1869  
*Palaeortyx gallica* MILNE-EDWARDS, 1869  
*Palaeortyx brevipes* MILNE-EDWARDS, 1869  
*Palaeortyx prisca* (MILNE-EDWARDS, 1869)  
 syn. *Palaeortyx intermedia* BALLMANN, 1966  
*Palaeortyx phasianoides* MILNE-EDWARDS, 1869  
*Palaeortyx media* MILNE-EDWARDS, 1869  
 (nomen nudum)

### O. Anseriformes

- F. Anatidae LEACH, 1820  
 Genus *Mionetta* LIVEZEY & MARTIN, 1988  
*Mionetta blanchardi* (MILNE-EDWARDS, 1863)  
*Mionetta consobrina* (MILNE-EDWARDS, 1867)  
*Mionetta natator* (MILNE-EDWARDS, 1867)  
 Genus *Cygnopterus* LAMBRECHT, 1931  
*Cygnopterus alphonsi* CHENEVAL, 1984

### O. Gaviiformes

- F. Gaviidae COUES, 1903  
 Genus *Colymboides* MILNE-EDWARDS, 1867  
*Colymboides minutus* MILNE-EDWARDS, 1867

- O. Procellariiformes  
F. Procellariidae LEACH, 1820  
Indeterminate genus  
Indeterminate genus *arvernensis* (MILNE-EDWARDS, 1871), syn. *Puffinus arvernensis* MILNE-EDWARDS, 1871, and *Plotornis arvernensis* (MILNE-EDWARDS, 1871) (CHENEVAL 1984; MAYR & SMITH 2012)
- O. Phoenicopteriformes  
F. Palaelodidae (STEJNEGER, 1885)  
Genus *Palaelodus* MILNE-EDWARDS, 1863  
*Palaelodus ambiguus* MILNE-EDWARDS, 1863, syn. *Ibidopodia palustris* MILNE-EDWARDS, 1871, syn. pro parte *Probalearica problematica* (MILNE-EDWARDS, 1868) (CHENEVAL & ESCUILLÉ 1992)  
*Palaelodus gracilipes* MILNE-EDWARDS, 1863, syn. *Palaelodus minutus* MILNE-EDWARDS, 1868 (CHENEVAL 1983)  
*Palaelodus crassipes* MILNE-EDWARDS, 1863  
Genus *Megapalaelodus* MILLER, 1944  
*Megapalaelodus goliath* (MILNE-EDWARDS, 1868)
- F. Phoenicopteridae BONAPARTE, 1831  
Genus *Phoenicopterus* LINNAEUS, 1758  
*Phoenicopterus croizeti* GERVAIS, 1852
- O. Ciconiiformes  
F. Ciconiidae GRAY, 1840  
Genus *Grallavis* CHENEVAL, 1984  
*Grallavis edwardsi* (LYDEKKER, 1891b)
- O. Pelecaniformes  
F. Threskiornithidae RICHMOND, 1917  
Genus *Gerandibis* DE PIETRI, 2013  
*Gerandibis pagana* (MILNE-EDWARDS, 1868) syn. *Milnea gracilis* LYDEKKER, 1891a (CHENEVAL 1984)
- F. Ardeidae LEACH, 1820  
Genus *Proardeola* HARRISON, 1979  
*Proardeola walkeri* HARRISON, 1979
- F. Pelecanidae RAFINESQUE, 1815  
Genus *Mioplecanus* CHENEVAL, 1984  
*Mioplecanus gracilis* (MILNE-EDWARDS, 1863)
- O. Suliformes  
F. Sulidae REICHENBACH, 1849  
cf. Genus *Emphesula* HARRISON, 1975  
cf. *Emphesula arvernensis* (MILNE-EDWARDS, 1867)
- F. Phalacrocoracidae REICHENBACH, 1850  
Genus *Oligocorax* LAMBRECHT, 1933  
*Oligocorax littoralis* (MILNE-EDWARDS, 1863)
- Genus *Nectornis* CHENEVAL, 1984  
*Nectornis miocaenus* (MILNE-EDWARDS, 1867)
- O. Accipitriformes  
F. Sagittariidae GRANDORI & GRANDORI, 1935  
Genus *Pelargopappus* STEJNEGER, 1885  
*Pelargopappus magnus* (MILNE-EDWARDS, 1868)
- F. Accipitridae VIEILLOT, 1816  
Genus *Aquilavus* LAMBRECHT, 1933  
*Aquilavus depredator* (MILNE-EDWARDS, 1871)  
*Aquilavus priscus* (MILNE-EDWARDS, 1863)  
Genus *Milvus* LACÉPÈDE, 1899  
*Milvus desperditus* MILNE-EDWARDS, 1871  
Genus *Promilio* WETMORE, 1958  
*Promilio incertus* (GAILLARD, 1939)  
Genus *Palaeohierax* MILNE-EDWARDS, 1871  
*Palaeohierax gervaisii* (MILNE-EDWARDS, 1863)
- O. Otidiformes  
F. Otididae RAFINESQUE, 1815 (?)  
Genus *Otis* LINNAEUS, 1758 (?)  
*Otis agilis* MILNE-EDWARDS, 1871 (nomen nudum)
- O. Gruiformes  
F. Rallidae VIGORS, 1825  
Genus *Palaeoaramides* LAMBRECHT, 1933  
*Palaeoaramides christyi* (MILNE-EDWARDS, 1869), syn. *Palaeoaramides eximius* (MILNE-EDWARDS, 1869)  
Genus *Paraortygometra* LAMBRECHT, 1933  
*Paraortygometra porzanoides* (MILNE-EDWARDS, 1869)
- F. Gruidae VIGORS, 1825  
Genus *Palaeogrus* PORTIS, 1884  
*Palaeogrus excelsus* (MILNE-EDWARDS, 1868)  
Genus *Probalearica* LAMBRECHT, 1933  
*Probalearica problematica* (MILNE-EDWARDS, 1868)
- O. Charadriiformes  
F. Haematopodidae BONAPARTE, 1838  
New material to be described (DE PIETRI *et al.* 2012)
- F. Recurvirostridae BONAPARTE, 1854  
Genus *Himantopus* BRISSON, 1760  
*Himantopus brevipes* MILNE-EDWARDS, 1871 (nomen nudum)
- F. Charadriidae VIGORS, 1825  
New material to be described (DE PIETRI *et al.* 2012)

Scolopaci sensu PATON *et al.* 2003

F. incertae sedis

Genus *Scolopacimilis* DE PIETRI & MAYR, 2012  
*Scolopacimilis lartetianus* (MILNE-EDWARDS, 1863)  
*Scolopacimilis* sp.

Genus *Becassius* DE PIETRI & MAYR, 2012  
*Becassius charadrioides* DE PIETRI & MAYR, 2012

F. Scolopacidae RAFINESQUE, 1815

Genus *Elorius* MILNE-EDWARDS, 1868  
*Elorius paludicola* MILNE-EDWARDS, 1868  
Genus ? *Elorius* MILNE-EDWARDS, 1868  
? *Elorius limosoides* DE PIETRI & MAYR, 2012  
Genus *Parvelorius* DE PIETRI & MAYR, 2012  
*Parvelorius gracilis* (MILNE-EDWARDS, 1868)  
Genus ? *Parvelorius* DE PIETRI & MAYR, 2012  
? *Parvelorius calidris* DE PIETRI & MAYR, 2012  
Genus and species indeterminate

Suborder Laromorphae DE PIETRI *et al.*, 2011a

F. Laricolidae DE PIETRI *et al.*, 2011a

Genus *Laricola* MLIKOVSKÝ, 2002  
*Laricola desnoyersii* (MILNE-EDWARDS, 1863)  
*Laricola elegans* (MILNE-EDWARDS, 1868)  
*Laricola totanoides* (MILNE-EDWARDS, 1868)  
*Laricola intermedia* DE PIETRI *et al.*, 2011a  
*Laricola robusta* DE PIETRI *et al.*, 2011a

Family incertae sedis

Genus *Sternalara* DE PIETRI *et al.*, 2011a  
*Sternalara minuta* DE PIETRI *et al.*, 2011a  
*Sternalara milneedwardsi* DE PIETRI *et al.*, 2011a

O. Pteroclidiformes

F. Pteroclididae BONAPARTE, 1831

Genus *Leptoganga* MOURER-CHAUVIRÉ, 1993  
*Leptoganga sepultus* (MILNE-EDWARDS, 1869)  
Genus *Gerandia* LAMBRECHT, 1933  
*Gerandia calcaria* (MILNE-EDWARDS, 1869)

O. Psittaciformes

F. Psittacidae ILLIGER, 1811

Genus *Archaeopsittacus* LAMBRECHT, 1933  
*Archaeopsittacus verreauxi* (MILNE-EDWARDS, 1871)

O. Strigiformes

F. Tytonidae RIDGWAY, 1914

Genus *Necrobyas* MILNE-EDWARDS, 1892  
*Necrobyas arvernensis* (MILNE-EDWARDS, 1863)

Genus *Prosybris* BRODKORB, 1970

*Prosybris antiqua* (MILNE-EDWARDS, 1863)

F. Strigidae VIGORS, 1825

Genus *Mioglaux* MLIKOVSKÝ, 1998  
*Mioglaux poirrieri* (MILNE-EDWARDS, 1863)

O. Apodiformes

F. Apodidae HARTERT, 1897

Genus *Procypseloides* HARRISON, 1984  
*Procypseloides ignotus* (MILNE-EDWARDS, 1871)

O. Coliiformes

F. Coliidae SWAINSON, 1837

Genus *Limnatornis* MILNE-EDWARDS, 1871  
*Limnatornis paludicola* MILNE-EDWARDS, 1871  
*Limnatornis archiaci* (MILNE-EDWARDS, 1871)

O. Trogoniformes

F. Trogonidae LESSON, 1828

Genus *Paratrogon* LAMBRECHT, 1933  
*Paratrogon gallicus* (MILNE-EDWARDS, 1871)

O. Coraciiformes

F. Coraciidae BATSCH, 1788

Genus *Miocoracias* nov. gen.  
*Miocoracias chenevali* nov. spec. (this paper)

O. Bucerotiformes

F. Phoeniculidae BONAPARTE, 1831

Genus *Phirriculus* MLIKOVSKÝ & GÖHLICH, 2000  
*Phirriculus pinicola* MLIKOVSKÝ & GÖHLICH, 2000

O. Piciformes

F. Picidae VIGORS, 1825

Genus *Piculoides* DE PIETRI *et al.*, 2011b  
*Piculoides saulcetensis* DE PIETRI *et al.*, 2011b

O. Passeriformes

F. Laniidae RAFINESQUE, 1815

Genus *Lanius* LINNAEUS, 1758  
*Lanius miocaenus* MILNE-EDWARDS, 1871

F. Ploceidae SUNDEVALL, 1836

Genus *Passer* BRISSON, 1760  
*Passer* sp.

F. Motacillidae HORSFIELD, 1821

Genus *Motacilla* LINNAEUS, 1758  
*Motacilla humata* MILNE-EDWARDS, 1871  
*Motacilla major* MILNE-EDWARDS, 1871

F. Fringillidae VIGORS, 1825

Genus *Loxia* LINNAEUS, 1758  
*Loxia* sp. 1  
*Loxia* sp. 2